

A SIMPLY CONNECTED NUMERICAL GODEAUX SURFACE WITH AMPLE CANONICAL CLASS

I. DOLGACHEV * and C. WERNER

1. Introduction

This paper is a follow-up of a recent paper of PierCarlo Craighero and Remo Gattazzo [CG] in which a new construction of a numerical Godeaux surface was given. Recall that a numerical Godeaux surface is a minimal surface V of general type with zero geometric genus p_g and $K_V^2 = 1$. The surface is obtained as a minimal resolution of a quintic surface in $\mathbb{P}^3$ with four simple elliptic singularities of type $z^2 + x^3 + y^6 = 0$. In this paper we show that the surface is simply connected and we also find its birational model as a double plane branched along an irreducible curve of degree 10 with five singular points of type $(3, 3)$ and one quadruple ordinary singular point. Previous examples of such a curve ([R],[St]) lead to classical Godeaux surfaces (with fundamental group of order 5). A reducible curve of this kind was constructed earlier by F. Oort and C. Peters [OP] and the second author [W]. The corresponding double cover is birationally equivalent to a numerical Godeaux surface with fundamental group $\mathbb{Z}/4\mathbb{Z}$ and $\mathbb{Z}/2\mathbb{Z}$, respectively.

Recall that $\{1\}, \mathbb{Z}/2\mathbb{Z}, \mathbb{Z}/3\mathbb{Z}, \mathbb{Z}/4\mathbb{Z}, \mathbb{Z}/5\mathbb{Z}$ are the only possible values for the torsion group of a numerical Godeaux surface. The first (and the only) construction of simply connected surfaces of general type with $p_g = 0$ was given by R. Barlow [B]. Although Barlow's surfaces are numerical Godeaux surfaces they are not isomorphic to the surfaces of Craighero-Gattazzo. In particular the Craighero-Gattazzo surfaces are the first examples of simply connected surfaces with $p_g = 0$ and ample canonical class. We prove this by showing that the surfaces contain no smooth rational curves with self-intersection (-2) ; however each Barlow surface contains four such curves. In [CL], F. Catanese and C. LeBrun prove that the Barlow surface can be deformed to a surface with ample canonical class. It is possible that Craighero-Gattazzo surfaces are deformation equivalent to Barlow surfaces.

Recently F. Catanese has announced a new construction of a family of simply connected numerical Godeaux surfaces. We do not know whether our surfaces belong to his family.

Finally the first author would like to thank E. Stagnaro for bringing to his attention the paper of Craighero and Gattazzo, and also M. Reid for useful discussions. Both authors thank J. Keum for his careful reading of the manuscript and many constructive comments.

2. The Quintics

Consider the following projective automorphism of $\mathbb{P}^3$:

$$\sigma : (X, Y, Z, T) \rightarrow (T, X, Y, Z).$$

* Research supported in part by a NSF grant.

Its order is four and its fixed points in $\mathbb{P}^3$ are

$$P_0 = (1, 1, 1, 1), P_1 = (1, i, -1, -i), P_2 = (1, -i, -1, i), Q_0 = (1, -1, 1, -1).$$

The set of fixed points of

$$\sigma^2 : (X, Y, Z, T) \rightarrow (Z, T, X, Y)$$

is equal to the union of two lines

$$r = \{X + Z = Y + T = 0\} = \langle P_1, P_2 \rangle, \quad r' : \{X - Z = Y - T = 0\} = \langle P_0, Q_0 \rangle.$$

We are looking for a quintic surface which is invariant with respect to σ and has simple elliptic singularities of degree 1 locally isomorphic to $z^2 + x^3 + y^6 = 0$ (tacnodal points) at the reference points

$$a_1 = (1, 0, 0, 0), \quad a_2 = (0, 1, 0, 0), \quad a_3 = (0, 0, 1, 0), \quad a_4 = (0, 0, 0, 1).$$

It is easy to check that any homogeneous polynomial $F_5(X, Y, Z, T)$ of degree 5 which is invariant with respect to σ and has degenerate critical points of multiplicity 2 at the points a_i must look like

$$(aT + bZ + Y)^2 X^3 + (aX + bT + Z)^2 Y^3 + (aY + bX + T)^2 Z^3 + (aZ + bY + X)^2 T^3 + (X^2 Y Z T + X Y^2 Z T + X Y Z^2 T + X Y Z T^2) c + (X^2 Y^2 Z + X^2 Y T^2 + X Z^2 T^2 + Y^2 Z^2 T) d + (X^2 Y^2 T + X^2 Z T^2 + X Y^2 Z^2 + Y Z^2 T^2) e + (X^2 Y Z^2 + X^2 Z^2 T + X Y^2 T^2 + Y^2 Z T^2) f.$$

One can show that for the values of the parameters

$$a = u^2, \quad b = -\frac{u^2 - u + 1}{2u^2 - 4u + 1}, \quad c = -\frac{34u^2 - 18u - 7}{29u^2 + 22u - 33},$$

$$d = \frac{7u^2 + 4u - 6}{3u - 2}, \quad e = -\frac{3u^2 + 6u - 8}{3u^2 + u - 2}, \quad f = -\frac{225u^2 - 156u - 10}{5u^2 + 212u - 163},$$

where $u^3 + u^2 - 1 = 0$, the surface has the required singularities (see [CG]). Also, the σ -invariance implies that F_5 vanishes at Q_0 and is identically zero along r .

Let S be the surface in $\mathbb{P}^3$ given by the equation $F_5 = 0$ and let $\pi : V \rightarrow S$ be a minimal resolution of singularities. Let $E_i = \pi^{-1}(a_i)$. By the adjunction formula,

$$K_V = \pi^*(H) - E_1 - E_2 - E_3 - E_4, \tag{2.1}$$

where H is a hyperplane section of the quintic. Since the points a_i are not coplanar,

$$p_g(V) = h^0(K_V) = 0.$$

Now one checks that the pencil of quadrics

$$\lambda YT + \mu XZ = 0 \tag{2.2}$$

defines the bicanonical linear system on V . It follows from the classification of algebraic surfaces that V is of general type. Noether's inequality together with $p_g(V) = 0$ imply that the irregularity $q(V)$ of V is also equal to zero.

We also see that the pencil $\lambda YT + \mu XZ = 0$ has four smooth base points, plus the singular points of the quintic. Thus the bicanonical system has no fixed part on the canonical model, therefore no fixed part on the minimal model. This implies that V is minimal.

We have

$$E_i^2 = -1, \quad E_i \cdot K_V = 1. \quad (2.3)$$

This gives

$$K_V^2 = 5 - 4 = 1,$$

thus V is a numerical Godeaux surface.

3. From quintic to double plane

In this section we shall find a double plane model of the surface V .

Let

$$R = \pi^{-1}(r) \quad (3.1)$$

be the inverse image of the line lying on S which is fixed by the involution σ^2 . This is a smooth rational curve on V . Since the line r does not contain the points a_i , we have

$$K_V \cdot R = 1, \quad R^2 = -3. \quad (3.2)$$

Consider the linear system

$$|\pi^*(H) - R|. \quad (3.3)$$

This is the pre-image of the pencil of plane quartic curves cut out on S by planes through the line r . We have

$$(\pi^*H - R) \cdot K_V = 4, \quad (\pi^*H - R)^2 = 0.$$

Thus $|\pi^*H - R|$ is a pencil of curves of genus 3.

The line r' intersects S at 5 points. Let

$$\Sigma = \{Q_0, Q_1, Q_2, Q_3, Q_4\}$$

be the set of the corresponding points on V . They are isolated fixed points of σ^2 on V . The point Q_0 is fixed by σ . The other two fixed points of σ on V lie on R .

Let $a : V' \rightarrow V$ be the blowing up of the points $Q_0, Q_1, \dots, Q_4$. The involution σ^2 acts on V' with the set of fixed points equal to the union of the curve

$$R' = a^{-1}(R) \quad (3.4)$$

and the curves

$$R_i = a^{-1}(Q_i), \quad i = 0, \dots, 4. \quad (3.5)$$

Let

$$p : V' \rightarrow F = V'/(\sigma^2) \quad (3.6)$$

be the canonical projection. The surface F is nonsingular.

Proposition 3.1. *F is a rational surface. The pencil $|\pi^*H - R|$ is the pre-image under $p : V' \rightarrow F$ of a pencil $|B|$ of elliptic curves on F .*

Proof. We have

$$K_{V'} = p^*(K_F) + \sum_{i=0}^4 R_i + R'.$$

This implies that the image of $p^*(2K_F)$ on V is equal to $2K_V - 2R$. But the linear system $|2K_V - 2R|$ is empty since no quadric from the pencil $\lambda YT + \mu XZ = 0$ contains the line r . This implies that $|2K_F| = \emptyset$, hence F is rational.

Each fiber in the pencil $|\pi^*H - R|$ is invariant with respect to σ^2 . Its general member intersects the ramification locus of the projection $p : V' \rightarrow F$ at four points lying on R , so the quotient $(\pi^*H - R)/\sigma^2$ is a curve of genus 1. Thus $|\pi^*H - R|$ defines an elliptic pencil $|B|$ on F .

Let us compute the topological Euler-Poincaré characteristic $e(F)$ of the surface F . By Noether's formula $12(1 - q + p_g) = K^2 + e$, we have $e(V) = 11$. Thus $e(V') = 16$ and the formula $e(V') = 2e(F) - e(R_0 + \dots + R_4 + R')$ gives

$$e(F) = 14. \tag{3.7}$$

Since $p_g(F) = q(F) = 0$, we must have $K_F^2 = -2$. Now observe that the elliptic curves E_i on V form two orbits with respect to σ^2 . Let $\bar{E}_1$ and $\bar{E}_2$ be the two elliptic curves on F representing these two orbits. Since each E_i does not intersect a general member of $|\pi^*H - R|$, the curves $\bar{E}_1, \bar{E}_2$ are contained in fibres of $|B|$. Also $\bar{E}_i^2 = -1$. Since for any minimal elliptic surface X we have $K_X^2 = 0$, a relative minimal model of F is an elliptic surface F' with $e(F') = 12$. This implies that F is obtained from F' by blowing up two points. Let B'_1 and B'_2 be the exceptional curves. The corresponding two elliptic fibres of F are the divisors $B_1 = \bar{E}_1 + B'_1, B_2 = \bar{E}_2 + B'_2$. They correspond to the plane sections $X + Z = 0$ and $Y + T = 0$ of the quintic surface.

We next consider the linear system $|3K_V - R|$.

Proposition 3.2. *$|3K_V - R|$ is a pencil of curves of genus 2, without fixed part.*

Proof. By Riemann-Roch, $\dim H^0(V, \mathcal{O}_V(3K_V)) = 4$. The tricanonical linear system defines a σ^2 -equivariant rational map from V to $\mathbb{P}^3$. Since σ^2 is identical on R , the image of R is contained in the fixed locus of σ^2 in $\mathbb{P}^3$. This fixed locus is the union of two skew lines or the union of a plane and a point. Since the preimage of this fixed locus on V is equal to R plus five isolated points, we see that the fixed locus in $\mathbf{P}^3$ cannot be a plane. Also R is not mapped to a point, thus the image of R is a line and hence the restriction of $|3K_V|$ to R is a pencil. By considering the exact sequence

$$0 \rightarrow \mathcal{O}_V(3K_V - R) \rightarrow \mathcal{O}_V(3K_V) \rightarrow \mathcal{O}_R(3K_V) \rightarrow 0$$

we obtain that $h^0(3K_V - R) = 2$. Thus $|3K_V - R|$ is a pencil.

It follows from (3.2) that

$$(3K_V - R)^2 = 9 - 6 - 3 = 0, \quad (3K_V - R) \cdot K_V = 3 - 1 = 2;$$

by the adjunction formula, the genus of the pencil is two.

We next show that this pencil has no fixed part. Assume there is a fixed part F , with

$$|3K_V - R| = |M| + F. \quad (3.8)$$

Intersecting both sides of (3.8) with K_V , we obtain $K_V \cdot M \leq 2$. The moving part $|M|$ must have genus two, so $M \cdot K_V = 1$ and $M^2 = 1$, or $M \cdot K_V = 2$ and $M^2 = 0$. Intersecting both sides of (3.8) with M gives

$$3K_V \cdot M - M \cdot R = M^2 + M \cdot F.$$

Since $|M|$ is obviously σ^2 -invariant, and σ^2 has isolated fixed points, each member of $|M|$ is σ^2 -invariant. Since its genus is 2, σ^2 has 2 or 6 fixed points on a general member. This gives $M \cdot R = 6$ or $M \cdot R = 2$. The first case immediately gives $M^2 = 0$ and $M \cdot F = F^2 = 0$. This contradicts the assumption that $F \neq 0$. Assume $M \cdot R = 2$. Since $M \cdot F > 0$, this is possible only if

$$M^2 = 0, \quad M \cdot F = 4, \quad F^2 = -8, \quad F \cdot K_V = 0, \quad F \cdot R = 4.$$

It is easy to see that this implies that F is the sum of four disjoint (-2) -curves each intersecting M with multiplicity 1. None of them is σ^2 -invariant (otherwise the intersection point with each member of $|M|$ is σ^2 -invariant). Thus σ acts transitively on the components of F , so that none of them contains a fixed point of σ^2 . However this contradicts the fact that $F \cdot R > 0$.

Lemma 3.3. *Let τ be an involution on a smooth surface X , and let W be a one-dimensional irreducible component of the locus of fixed points of τ . Let $C \neq W$ be an irreducible τ -invariant curve on X which intersects W at a point x . Then either $\text{mult}_x(C, W) = 1$ or x is a singular point of C . If x is an ordinary node of C , then $\text{mult}_x(C, W) = 2$. If x is an ordinary cusp point of C then $\text{mult}_x(C, W) = 3$.*

Proof. Since τ acts identically on W , it acts identically on the tangent space $T(W)_x$. If x is a nonsingular point of C and C is tangent to W at x , then τ acts on the tangent space $T(C)_x = T(W)_x$ identically. This of course implies that τ acts identically on C . However this contradicts the fact that the locus of fixed points of τ is smooth. Assume that C has a node; then τ switches the tangent directions at the two branches, so each branch intersects W with multiplicity 1. If C has a cusp at x , then the cuspidal tangent direction coincides with the tangent direction of W at x , so $\text{mult}_x(C, W) = 3$.

Lemma 3.4. *Let σ be an involution on a nonsingular surface X . Let C be a σ -invariant curve on X on which σ does not act identically. Assume that C passes through k isolated fixed points p_i of σ on X . Then*

$$C^2 \equiv \sum_{i=1}^k (\text{mult}_{p_i} C)^2 \pmod{2}.$$

Proof. Let $X' \rightarrow X$ be the blowing up of isolated fixed points of σ and $\bar{C}$ be its proper inverse transform. Let $\bar{X} = X'/(\sigma)$. This is a nonsingular surface and $\bar{C}$ is equal to the pre-image of a curve on it not contained in the branched locus of the projection $X' \rightarrow \bar{X}$. Therefore $\bar{C}^2 = C^2 - \sum_{i=1}^k (\text{mult}_{p_i} C)^2$ is even.

Lemma 3.5. *Let $\Sigma = \{Q_0, \dots, Q_4\}$ be the set of isolated fixed points of σ^2 . Let D be a member of the pencil $|3K_V - R|$ such that $\Sigma \cap D \neq \emptyset$. Then one of the following cases occurs:*

- (i) $\#\Sigma \cap D = 1, D = A_1 + A_2$ is the union of two irreducible curves of arithmetic genus 1 with $A_1 \cap A_2 = \{Q_i\}$;
- (ii) $\#\Sigma \cap D = 1, D = 2A + N_1 + N_2$, where A is an irreducible curve of arithmetic genus 1 passing through $Q_i \in \Sigma$ and $N_1 = \sigma^2(N_2)$ are two disjoint (-2) -curves intersecting A with multiplicity 1;
- (iii) $\#\Sigma \cap D = 2, D = A_1 + A_2 + N$ where A_1, A_2 are two disjoint irreducible curves of arithmetic genus 1, each passing through one of the two points of $\Sigma \cap D$, and N is a (-2) -curve intersecting A_1 and A_2 each with multiplicity one at these points.

Proof. Since $D \cdot K_V = (3K_V - R) \cdot K_V = 2$, we can write $D = A + N$, where N is the union of (-2) -curves and $K_V \cdot A = 2$. The latter equality shows that A has at most two irreducible components. For any component C of N , we have $C \cdot (3K_V - R) = -C \cdot R$, therefore $C \cdot R = 0$ and $C \cdot (3K_V - R) = 0$. Thus

$$R \cdot A = R \cdot D = R \cdot (3K_V - R) = 6.$$

If A is an irreducible curve of arithmetic genus 2, then $A^2 = 0$ and $D = A$. By Lemma 3.3, D has either six fixed points of σ^2 on R , or D has a node along R and four additional fixed points on R , or two nodes and two additional fixed points, or a cusp along R and three additional fixed points from R , or 2 cusps and no fixed points on R . Since D has at least one fixed point away from R it is easy to see that σ^2 acts on the normalization of D with an odd number of fixed points. This obviously contradicts the Hurwitz formula.

A similar argument shows that D cannot be irreducible and reduced of arithmetic genus less than 2.

Thus A is reducible or multiple. Observe that D , and hence A and N , are σ^2 -invariant. Since a nonsingular rational component of A has at most two fixed points, the equality $A \cdot R = 6$ shows that A cannot consist of two nonsingular rational components. Also if C is a nonsingular rational component of A , then the remaining component C' must satisfy $K_V \cdot C' = 1$ and hence must be of arithmetic genus 1. Since $R \cdot (C + C') = 6$ the rational component C must have two distinct intersection points with R (Lemma 3.3) and no more fixed points of σ . Since $C^2 = -3$ is odd this contradicts Lemma 3.4.

Thus each irreducible component of A is of arithmetic genus 1. Notice that since $A \cdot R = 6$, we must have two irreducible components A_1 and A_2 which of course may coincide. Obviously, each A_i is σ^2 -invariant. By Lemma 3.4 each A_i contains an odd number of nonsingular points from Σ . Since $(A_1 + A_2) \cdot R = 6$ and $A_i \cdot R \leq 4$, we obtain that each A_i contains exactly one nonsingular point from the set Σ .

Case 1: $A_1 \neq A_2, A_1 \cdot A_2 \neq 0$. In this case A_1 and A_2 have a common point from Σ and $A_1 \cdot A_2 = 1$. The formulae $0 = D \cdot A_i = (A_1 + A_2 + N) \cdot A_i = -1 + 1 + N \cdot A_i$ imply that $N = 0$. This is case (i).

Case 2: $A_1 \neq A_2, A_1 \cdot A_2 = 0$. As above we obtain that each A_i intersects N at one point. This point must be one of the points Q_i . Since $0 = (A_1 + A_2 + D_2)^2 = 2 + N^2$, we see that $N^2 = -2$. Let $N_1, \dots, N_k$ be the irreducible components of N such that N_1 intersects A_1 and N_k intersects A_2 . Clearly, N_1 and N_k are σ^2 -invariant. If $k > 1$, i.e. $N_1 \neq N_k$, each N_i contains an additional point from Σ . We know that the image of Σ in S lies on the line r' . Since $N \cdot K_V = 0$, N_1 and N_k must be components of a member of the pencil $|2K_V|$; this system corresponds to quadrics in $\mathbb{P}^3$, which can intersect the line r' in at most two points. Therefore $N_1 \cup N_k$ must have at most two points from Σ . This contradiction shows that $N_1 = N_k$ and hence N is a single (-2) -curve. This gives us case (iii).

Case 3: $A_1 = A_2$ so that $3K_V - R = 2A_1 + N$. Intersecting both sides with K_V we get $A_1 \cdot K_V = 1, A_1^2 = -1$. The formulae

$$0 = D \cdot A_1 = (2A_1 + N) \cdot A_1 = -2 + N \cdot A_1,$$

$$D^2 = -4 + 4A_1 \cdot N + N^2 = 0$$

show that $N \cdot A_1 = 2, N^2 = -4$. The last equality shows that N consists of at least two irreducible components. It cannot consist of two components N_1 and N_2 with $N_1 \cdot N_2 = 1$. Indeed, write $N = mN_1 + nN_2$ to get

$$N^2 = -2m^2 - 2n^2 + 2mn = -2(m - n)^2 - 2mn.$$

It is easy to see that N^2 cannot be equal to -4 . So, $N = N_1 + N_2$, where $N_1 \cdot N_2 = 0$ as in case (ii), or N contains more than two irreducible components. It remains to exclude this case. Observe that D is not σ -invariant since otherwise A_1 is σ -invariant and hence has two fixed points of σ (it is an easy fact that which follows from the Hurwitz formula that any involution of order 4 on an irreducible curve of arithmetic genus 1 has 2 fixed points). Let $D_1, \dots, D_k$ be the divisors from $|3K_V - R|$ which contain points of Σ and let r_i be the number of irreducible components of D_i . Since $b_2(V) = \text{rk Pic}(V) = 9$, we have $\sum_i (r_i - 1) \leq 7$. If our N contains more than three components, D and $\sigma(D)$ will contribute at least $4 + 4$ to this sum which is too much. Since D is σ^2 -invariant, N must be supported on the nodal cycle $N_1 + N_2 + N_3$ of type A_3 with N_1 and N_3 intersecting A . Moreover N_2 must be σ^2 -invariant and hence contains two points from Σ . Thus D contains three points from Σ , and $\sigma(D)$ contains three more points from Σ which is too many.

This proves the assertion.

Corollary 3.6. *In the notation of the previous lemma, let $D_i, i = 1, \dots, k$, be the set of divisors from $|3K_V - R|$ such that $D_i \cap \Sigma \neq \emptyset$. Then one of the following cases occurs:*

- (i) $k = 5$, each D_i is of type (i);
- (ii) $k = 5$, D_4 and $D_5 = \sigma(D_4)$ are of type (ii), D_1, D_2, D_3 are of type (i);

(iii) $k = 3$, D_2 and $D_3 = \sigma(D_1)$ are of type (iii), D_1 is of type (i).

Moreover, the unique σ -invariant divisor, say D_1 , is of type (i).

Proof. Since Σ consists of five points, and each divisor D_i contains at most two points from Σ , we have $5 \geq k \geq 3$. Assume $k = 5$. Then each D_i is of type (i) or (ii). Let r_i be the number of irreducible components of D_i . As we already noticed in the previous proof, $\sum_i (r_i - 1) \leq 7$. This implies that we have at most two divisors D_i of type (ii). If there is only one D_i of type (ii), then it is necessary σ -invariant. But then each of the rational components of D_i is σ^2 -invariant. Since it does not intersect R and does not contain points from Σ , we get a contradiction. This gives us two cases (i) and (ii) with $k = 5$.

Assume $k = 4$. Then exactly one of the divisors D_i must contain two points from Σ , and hence it must be of type (iii) and σ -invariant. But then its rational component must be invariant and contain two isolated fixed points of σ . This is impossible since we have only one such point. So this case does not occur.

Assume $k = 3$. Then we must have two divisors of type (iii). The third divisor cannot be of type (ii) since otherwise it is σ -invariant, and we have already seen that this is impossible.

Theorem 3.7. *V is isomorphic to a nonsingular minimal model of the double cover of $\mathbb{P}^2$. The branch curve is an irreducible curve W of degree 10 with a point q of multiplicity 4 and 10 triple points p_i, p'_i , $i = 0, \dots, 4$, where p'_i is infinitely near to p_i . The points $p_0, \dots, p_4, q$ do not lie on a conic. Also there exists a pencil $|\bar{B}|$ of elliptic curves of degree 4 with double points at p_0 and p'_0 and simple points at other triple points of W .*

Proof. We shall employ the notation from the proof of Proposition 3.1. Let B be the pencil of elliptic curves on $F = V'/(\sigma^2)$ whose general fibre is the quotient of the general fibre of $|\pi^*(H) - R|$ by (σ^2) . Let $|D|$ be the pencil of rational curves on F whose general fibre is the quotient of the general fibre of $|3K_V - R|$ by (σ^2) . It follows from (3.2) that $(3K_V - R) \cdot (\pi^*(H) - R) = 8$, hence

$$B \cdot D = \frac{1}{2}(3K_V - R) \cdot (\pi^*(H) - R) = 4.$$

Let $B_1 = \bar{E}_1 + B'_1 \in |B|$ be one of the two non-minimal fibres of the elliptic fibration defined by $|B|$. The curve B'_1 is the exceptional curve of the first kind and the pre-image of $\bar{E}_1$ is the union of two elliptic curves E_1 and E_3 on V . Since $(E_1 + E_3) \cdot (3K_V - R) = 6$, we get

$$D \cdot \bar{E}_1 = 3, \quad D \cdot \bar{B}'_1 = 1.$$

Thus B'_1 is a section of the pencil $|D|$.

Now consider the divisors $D_i, i = 1, \dots, k$, from $|3K_V - R|$ which contain a point from the set Σ of isolated fixed points of σ^2 . Lemma 3.5 and Corollary 3.6 describe these divisors.

Let $\bar{D}_i$ denote the corresponding divisor in $|D|$. If $D_i = A_1 + A_2$ is of type (i), then

$$\bar{D}_i = \bar{A}_1^{(i)} + \bar{A}_2^{(i)} + \bar{R}^{(i)},$$

where $\bar{A}_1^{(i)}, \bar{A}_2^{(i)}$ are exceptional curves of the first kind, and $\bar{R}^{(i)}$ is the image in F of $a^{-1}(Q) \in V'$, where $Q = D_i \cap \Sigma$.

If $D_i = 2A + N_1 + N_2$ is of type (ii), then similarly we have


$$\bar{D}_i = 2\bar{A} + \bar{N}^{(i)} + \bar{R}^{(i)}.$$

where $\bar{R}^{(i)}$ is the image of $a^{-1}(D_i \cap \Sigma)$ and $\bar{N}^{(i)}$ is the image of the orbit of (-2) -curves $N_1 + N_2$ on F .

Finally if D_i is of type (iii),

$$\bar{D}_i = \bar{A}_1^{(i)} + \bar{A}_2^{(i)} + \bar{R}_1^{(i)} + \bar{R}_2^{(i)} + \bar{N}^{(i)},$$

with the similar notation.



Reducible fibres of $|3K-R|$

Consider the section $\bar{B}'_1$. It intersects each divisor $\bar{D}_i$ simply at one point. Without loss of generality we may assume that this point does not lie on $\bar{A}_1^{(i)}$. Also we observe that $\bar{B}'_1$ does not intersect the the curve $\bar{R}^{(i)}$ (where $\bar{R}^i = \bar{R}_1^i + \bar{R}_2^i$ in case D_i is of type (iii)). This is because the planes $X + Z = 0$ and $Y + T = 0$ do not contain points from Σ .

For each $i = 1, \dots, k$, let $\mathcal{E}_i$ be the divisor chosen as follows. If D_i is of type (i) or (ii),

$$\mathcal{E}_i = \bar{A}_1^{(i)} + \bar{R}^{(i)},$$

if D_i is of type (iii),

$$\mathcal{E}_i = \bar{A}_1^{(i)} + \bar{R}_1^{(i)} + \bar{R}_2^{(i)} + \bar{N}^{(i)},$$

It is easy to see that starting from the curve $\bar{A}_1^{(i)}$ we can blow down the divisor $\mathcal{E}_i$ to a nonsingular point. Using Corollary 3.6, it is easy to check that the union of the divisors $\mathcal{E}_i, i = 1, \dots, k$, is an exceptional divisor with 10 irreducible components which can be blown down to k distinct nonsingular points. Let $\alpha : F \rightarrow \bar{F}$ be its blowing down. It follows from (3.7) that $e(\bar{F}) = 4$, hence $\bar{F}$ is a minimal ruled surface. Since each $\mathcal{E}_i$ is a part of a member of the pencil $|D|$, the image of $|D|$ on $\bar{F}$ defines a ruling

$$\alpha_1 : \bar{F} \rightarrow \mathbb{P}^1.$$

It is clear that the curve B'_1 does not intersect $\mathcal{E}_i$ if D_i is of type (i) or (ii). We shall prove a little later that case (iii) does not occur. Let us continue the proof accepting this fact. The image of B'_1 is a curve L on $\bar{F}$ with $L^2 = B_1'^2 = -1$. This implies that $\bar{F}$ is the minimal ruled surface $\mathbf{F}_1$. We can blow down the curve L to arrive at $\mathbb{P}^2$.

Since we have blown down all the curves $\bar{R}_j^{(i)}$, our surface V is birationally isomorphic to the double plane branched along the curve W equal to the image of the “line” R . The image $\bar{R}$ of R on F has self-intersection -6 . It intersects each $\mathcal{E}_i$ at the curve $\bar{A}_1^{(i)}$ with multiplicity 3 and intersects the curve L with multiplicity 4. When we blow down $\mathcal{E}_i$ we increase the self-intersection by 18. When we blow down L it increases by 16. Using Corollary 3.6 we see that in any case we increase $\bar{R}^2$ by 106, and hence $W^2 = 100$, i.e. W is a curve of degree 10. It has a quadruple point at the image of L . It has a triple point with one infinitely near triple point at the image of $\mathcal{E}_i$. Applying Corollary 3.6 we get the assertion about the degree and the singularities of the branch curve W .

It remains to check the assertion about the elliptic pencil. As the reader can already guess this must be the image of the pencil $|B|$. First of all, we have already noticed that $B \cdot D = 4$. This means that the image of $|B|$ on $\mathbf{F}_1$ intersects the ruling with multiplicity 4. This implies that the image $\bar{B}$ of B on $\mathbb{P}^2$ is a pencil of elliptic quartics. Let D_1 be the unique σ -invariant divisor among the divisors $D_1, \dots, D_k$. We know from Corollary 3.6 that it is of type (i). A general member B of the pencil $|\pi^*(H) - R|$ intersects each $D_1 = A_1 + A_2 \in |3K_V - R|$ with multiplicity 4. Since $\sigma(A_1) = A_2$ and $\sigma(\pi^*(H) - R) = \pi^*(H) - R$ this implies that $(\pi^*(H) - R) \cdot A_1 = (\pi^*(H) - R) \cdot A_2 = 4$, and hence

$$(B'_1 + \bar{E}_1) \cdot \bar{A}_1 = (B'_1 + \bar{E}_1) \cdot \bar{A}_2 = \bar{E}_1 \cdot \bar{A}_1 = 2.$$

Thus when we blow down $\mathcal{E}_1$ the image of B in $\mathbb{P}^2$ acquires a double point at p_0 and a double point at p'_0 . There are no more singular points for a general member of $|\bar{B}|$. So B intersects each $\mathcal{E}_i$ with multiplicity 1. This gives us the assertion about the pencil $|\bar{B}|$. The assertion about the conic follows easily from computing the canonical class of the double plane (see the proof of Theorem 4.1).

It remains to pay our debt by checking that case (iii) in Corollary 3.6 does not occur. Suppose D_i is of type (iii); we claim that B'_1 does not intersect $\mathcal{E}_i$ in this case. Suppose it does. Then we still can blow down the divisors $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ to arrive at a minimal ruled surface $\bar{F}$. The image $\bar{B}'_1$ of B'_1 on $\bar{F}$ is a section with positive self-intersection. Its divisor

class on $\bar{F}$ is equal to $nf + s$, $n \neq 0$, where f is the class of the fibre and s is the class of the exceptional section. Recall that $R' = R/(\sigma^2)$ is a multi-section of the pencil $|D|$ of degree 6. So the image $\bar{R}$ of R' on $\bar{F}$ intersects f with multiplicity 6. This gives

$$\bar{B}'_1 \cdot \bar{R} = (nf + s) \cdot \bar{R} \geq 6n \geq 6.$$

However B'_1 is the image on F of a component of a divisor from $|\pi^*(H) - R|$ which intersects R with multiplicity 4. This shows that $\bar{B}'_1 \cdot \bar{R} \leq 4$ and gives us a contradiction. So, even in the case when D_i is of type (iii) we can blow down the exceptional divisors $\mathcal{E}_i$ and B'_1 to obtain the plane model of F . The image of the pencil $|B|$ is again a pencil of quartics. Thus the self-intersection of B must decrease by 16. When we blow-down $\mathcal{E}_1$ and B'_1 we decrease it by 8. Thus $\mathcal{E}_2$ and $\mathcal{E}_3$ must contribute the remaining 8. Since the general member of $\bar{B}$ has only two singular points which give a tacnode at the image of $\mathcal{E}_1$, we see that $B \cdot \mathcal{E}_2 = B \cdot \mathcal{E}_3 = 1$. We also know that B does not intersect the components $R_j^{(i)}$. This easily gives that B^2 decreases by at most $3 + 3 = 6$, a contradiction. Thus, all the assertions of the theorem have been checked, and we obtain that case (iii) in Corollary 3.6 does not occur.

Remark 3.8. Recall that the elliptic pencil $|B|$ on F contains two reducible members $\bar{E}_1 + B'_1$ and $\bar{E}_2 + B'_2$. The image of $\bar{E}_1$ is a quartic from the pencil $|\bar{B}|$ passing through q . The curve B'_1 is of course blown down to q . The image of $\bar{E}_2$, the orbit of E_2 and E_4 , is a cubic curve which passes simply through the points $p_i, p'_i, i = 0, \dots, 4$. The image of B'_2 is the line passing through p_0 and p'_0 . One can make a standard (degenerate) Cremona transformation with center at the points p_0, p'_0, q . It transforms the branch curve R into itself, and replaces the curve L with the line l through (p_0, p'_0) . This is induced by the involution $\bar{\sigma}$.

Remark 3.9. We shall later show that also the case (ii) in Corollary 3.7 does not occur for the surfaces which we are considering. It is quite possible that cases (ii) and (iii) occur for some genus 2 pencil on a numerical Godeaux surface. Observe that the case (iii) corresponds to the degeneration when, say $p_1 = p_2$ and $p_3 = p_4$. The case (ii) corresponds to the degeneration when the points q, p_i, p'_i become collinear. .

Remark 3.10. We can also realize V as a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ as follows. Consider the linear system $|4K_V - R|$; we have

$$(4K_V - R)^2 = 5, (4K_V - R) \cdot K_V = 3, (4K_V - R) \cdot (3K_V - R) = 2.$$

Thus $|4K_V - R|$ is a linear system of genus five curves. One checks that this is a pencil with five base points, which are exactly the points $Q_0, Q_1, \dots, Q_4$ fixed by the involution σ^2 . We know that the members of $|3K_V - R|$ passing through these points contain pairs of elliptic curves $A_i + A'_i$.

Thus these two pencils define a double cover $V \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$. This map blows up the base points of $|4K_V - R|$ to rulings of $\mathbb{P}^1 \times \mathbb{P}^1$ and contracts the elliptic curves to pairs of points on these fibres (for details, see [R]). In case (ii) of the Lemma, we have $A_i = A'_i$ for a member of $|3K_V - R|$; in this case the pair of points on the corresponding fibre are infinitely near.

We have

$$(3K_V - R) \cdot R = 6, \quad (4K_V - R) \cdot R = 7,$$

thus the rational curve R is mapped to a curve of bidegree $(6, 7)$ with pairs of triple points on the five special fibres of $|3K_V - R|$. This curve on $\mathbb{P}^1 \times \mathbb{P}^1$ can then be birationally transformed into a degree ten plane curve as in the theorem.

4. From double plane to quintic

One can reverse the previous construction.

Theorem 4.1. *Assume there exists an irreducible plane curve W of degree 10 with an point q of multiplicity 4 and 10 triple singular points $p_i, p'_i, i = 0, \dots, 4$, where p'_i is infinitely near to p_i . We assume that the points $p_0, \dots, p_4, q$ do not lie on a conic. Also assume that there exists a pencil D of elliptic curves of degree 4 with double points at p_0 and p'_0 and simple points at $p_i, p'_i, i \neq 0$. Let V' be a double cover of $\mathbb{P}^2$ branched over W . Then its nonsingular minimal model V is a numerical Godeaux surface. Let E be the pre-image of the elliptic pencil on V and R be the proper inverse transform of the curve W ; then the linear system $|E + R|$ maps V onto a normal quintic in $\mathbb{P}^3$ with 4 simple elliptic singular points of degree 1. The singular points of the quintic are the images of the inverse transforms of the quartic Q from the pencil which passes through q and the cubic curve C which together with the line $l = \langle p_0, p'_0 \rangle$ forms a member of the pencil of elliptic curves. If additionally, W is invariant with respect to the standard Cremona transformation with center p_0, p'_0, q , then the quintic is invariant with respect to a projective transformation of order 4.*

Proof. It is rather standard. Let $\pi : F \rightarrow \mathbb{P}^2$ be the blowing-up of points $q, p_i, p'_i, i = 0, \dots, 4$. Then

$$K_F = -3h + Z + \sum_{i=0}^4 (Z_i + 2Z'_i), \quad (4.1)$$

where h is the pre-image of a line, $Z_i + Z'_i = \pi^{-1}(p_i)$, $Z'_i = \pi^{-1}(p'_i)$, $Z = \pi^{-1}(q)$. The proper transform of W is

$$\bar{W} = 10h - 4Z - \sum_{i=0}^4 (3Z_i + 6Z'_i). \quad (4.2)$$

We take the double cover $p : V' \rightarrow F$ branched along the nonsingular divisor

$$W' = \bar{W} + \sum_{i=0}^4 Z_i. \quad (4.3)$$

Then, we have

$$K_{V'} = p^*(K_F + \frac{1}{2}W') = 2p^*(h) - p^*(Z) - \sum_{i=0}^4 p^*(Z'_i).$$

It follows that

$$p_*(\mathcal{O}_{V'}(K_{V'})) = \mathcal{O}_F(K_F + \frac{1}{2}W) \oplus \mathcal{O}_F(K_F) = \mathcal{O}_F(2h - Z - \sum_{i=0}^4 Z'_i) \oplus \mathcal{O}_F(K_F).$$

Since $h^0(2h - Z - \sum_{i=0}^4 Z'_i) = 0$ (otherwise there would be a conic through the points q, p_i), we obtain

$$h^0(K_{V'}) = h^0(p_*(\mathcal{O}_{V'}(K_{V'}))) = 0.$$

By Riemann-Roch, $h^1(2h - Z - \sum_{i=0}^4 Z'_i) = 0$, so we obtain

$$h^1(K_{V'}) = h^1(p_*(\mathcal{O}_{V'}(K_{V'}))) = 0.$$

On the other hand,

$$|2K_{V'}| = |p^*(4h - 2Z - 2\sum_{i=0}^4 Z'_i)| \neq \emptyset,$$

since it contains the positive divisors equal to the sum of the curves Z_i and the proper transform of a quartic curve C_4 from the pencil of quartics passing simply through the points p_i, p'_i and with multiplicity 2 at q . The curves $\bar{A}_i = p^*(Z'_i)$ are elliptic curves with self-intersection -2 . The curves $R_i = p^*(Z_i)$ are exceptional curves of the first kind. Blowing down the curves R_i we get a surface V with

$$K_V^2 = K_{V'}^2 + 5 = 1.$$

Together with the property $h^0(2K_V) \neq 0$, this shows that V is a surface of general type with

$$p_g = q = 0.$$

It is also minimal. In fact any exceptional curve of the first kind intersects $2K_V$ negatively, hence must be a component of the image on V of the proper transform of any quartic curve C_4 from above. It is easy to see that the pencil of such quartics does not have base components. Thus V is a numerical Godeaux surface.

Let $C_1 \sim h - Z_0 - 2Z'_0$ be the proper transform on F of the line l , and $C_2 = Z$. The pre-image on F of the elliptic pencil formed by the quartic curves is equal to

$$4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i);$$

we have

$$4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i) \sim C_1 + [3h - \sum_{i=0}^4 (Z_i + 2Z'_i)]$$

and

$$4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i) \sim C_2 + [4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i) - Z].$$

The curves $B_1 \in |3h - \sum_{i=0}^4 (Z_i + 2Z'_i)|$ and $B_2 \in |4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i) - Z|$ are elliptic curve on F of self-intersection -1 which are disjoint from the branch divisor W' . They are equal to the proper transforms of the cubic curve C and the quartic Q , respectively. Under the double cover $p : V' \rightarrow F$ they are split into the disjoint sum of two elliptic curves with self-intersection -1 , $p^*(B_1) = E'_1 + E'_2$ and $p^*(B_2) = E'_3 + E'_4$. We have

$$\begin{aligned} K_{V'} + E'_1 + E'_2 + E'_3 + E'_4 &\sim [p^*(2h - Z - \sum_0^4 Z'_i)] + \\ &+ [p^*(4h - Z - 2Z_0 - 4Z'_0 - \sum_1^4 (Z_i + 2Z'_i))] + [p^*(3h - \sum_0^4 (Z_i + 2Z'_i))] \\ &= 9p^*(h) - 2p^*(Z) - 3p^*(Z_0) - 7p^*(Z'_0) - 2 \sum_{i=1}^4 p^*(Z_i) - 5 \sum_{i=1}^4 p^*(Z'_i). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} p^*(4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i)) + \frac{1}{2}p^*(\bar{W} + \sum_{i=0}^4 Z_i) &= \\ = p^*(4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i)) + p^*(5h - 2Z - \sum_{i=0}^4 (Z_i + 3Z'_i)). \end{aligned}$$

Comparing, we get that

$$K_{V'} + E'_1 + E'_2 + E'_3 + E'_4 \sim p^*(D') + \frac{1}{2}p^*(\bar{W} + \sum_{i=0}^4 Z_i),$$

where $|D'| = |p^*(4h - 2Z_0 - 4Z'_0 - \sum_{i=1}^4 (Z_i + 2Z'_i))|$ is the pencil of curves originating from the pencil of quartic curves. Blowing down the exceptional curves p^*Z_i , we get

$$K_V + E_1 + E_2 + E_3 + E_4 \sim R + D,$$

where E_i, D are the images of E'_i, D' , and R is the reduced pre-image of $\bar{W}$. Clearly, R is a nonsingular model of our branch curve of degree 10. Since the genus of the latter is zero, R must be isomorphic to $\mathbb{P}^1$. Let us denote the linear system $|R + D|$ by $|H|$ and prove that $|H|$ maps V birationally onto a normal quintic in $\mathbb{P}^3$.

We immediately check that $D \cdot R = 4$, hence $|D|$ is a pencil of curves of genus 3. Since $D \cdot D = 0$, we obtain

$$D \cdot K_V = 4.$$

From this it follows that

$$5 = K_V^2 + K_V \cdot (E_1 + \dots + E_4) = K_V \cdot D + K_V \cdot R = 4 + K_V \cdot R,$$

hence

$$K_V \cdot R = 1, \quad R^2 = -2 - 1 = -3.$$

This immediately implies

$$H^2 = (D + R)^2 = 5, \quad H \cdot K_V = 5$$

Since we know that $p_g(V) = q(V) = 0$, by Riemann-Roch

$$h^0(H) = \frac{1}{2}(H^2 - K_V \cdot H) + 1 + h^1(H) = 1 + h^1(H) = 1 + h^1(K_V - H).$$

Now the exact sequence

$$0 \rightarrow \mathcal{O}_V(-E_1 - E_2 - E_3 - E_4) \rightarrow \mathcal{O}_V \rightarrow \mathcal{O}_{E_1+E_2+E_3+E_4} \rightarrow 0$$

together with the isomorphism $\mathcal{O}_V(-E_1 - E_2 - E_3 - E_4) \cong \mathcal{O}_V(K_V - H)$ gives

$$h^1(H) = h^1(-E_1 - E_2 - E_3 - E_4) = h^0(\mathcal{O}_{E_1+E_2+E_3+E_4}) - h^0(\mathcal{O}_V) = 3.$$

Thus

$$\dim |H| = h^0(H) - 1 = 3.$$

The complete linear system does not have base points. In fact, since $H = D + R$, where $|D|$ is a base-point-free pencil, any base point of H belongs to R . But $H \cdot R = 1$ and the exact sequence

$$0 \rightarrow \mathcal{O}_V(D) \rightarrow \mathcal{O}_V(H) \rightarrow \mathcal{O}_R(H) \rightarrow 0$$

shows easily that $|H|$ cuts out a pencil on R . This checks that there are no base points on R .

Now the linear system $|H|$ defines a regular map $f : V \rightarrow \mathbb{P}^3$. Since $H \cdot E_i = 0$, it blows down the curves E_i . Since $H^2 = 5$ is a prime number, f must be of degree 1 with the image a quintic surface with 4 elliptic singular points of degree 1. Since a nonsingular member of H is of genus $g = 1 + \frac{1}{2}(H^2 + H \cdot K_V) = 6$, and this coincides with the arithmetic genus of a plane section of a quintic surface, the image of f is a normal surface. So, all the assertions of the theorem are proven.

5. The fundamental group

We shall start with the following

Lemma 5.1. *Assume V contains a (-2) -curve C . Then C is a component of one of the members of the pencil $|3K_V - R|$ which passes through a point $Q_i, i \neq 0$. Moreover, its image in the quintic model is a conic which passes through two of the four singular points a_i .*

Proof. Suppose C is a (-2) -curve. Since $(3K_V - R) \cdot C = -R \cdot C$ and $C \neq R$, we get that $C \cdot R = 0$ and C is contained in a member of $|3K_V - R|$. If C does not contain one of the points Q_i , then the involution σ^2 has no fixed points on C , and the orbit of C under the group (σ) consists of 4 disjoint (-2) -curves. Using the argument from the proof of Corollary 3.6, we find that there are too many irreducible components contained in reducible fibres of $|3K_V - R|$.

Thus C must belong to one of the divisors D_i from Corollary 3.6. As we have already noticed in the proof of Theorem 3.7, the case (iii) of Corollary 3.6 does not occur for our surfaces. Thus there are four of the curves C forming an orbit with respect to (σ) . Each (σ^2) -orbit is contained in one reducible fibre of $|3K_V - R|$.

Let $D = 2A + C + \sigma^2(C)$ be one of the reducible members of the pencil $|3K_V - R|$ which contains a (-2) -curve C as in case (ii) of Lemma 3.5, and let E_i be one of the four elliptic curves blown up from the singular points of the quintic. Then $E_i \cdot (3K_V - R) = 3$. Hence $3 = E_i \cdot (2A + C + \sigma^2(C)) = 2(E_i \cdot A) + E_i \cdot C + E_i \cdot \sigma^2(C)$. It follows from the proof of Theorem 3.7 that $E_i \cdot A \neq 0$. This implies that $E_i \cdot A = 1$ and E_i intersects $C + \sigma^2(C)$ at one point. Now repeating this argument for the member $\sigma(D)$ of $|3K_V - R|$ we find that E_i intersects one more (-2) -curve $\sigma(C)$ (or $\sigma^3(C)$). Thus each E_i intersects exactly two (-2) -curves. This implies that each (-2) -curve C intersects two elliptic curves E_i each with multiplicity 1. Now

$$0 = K_V \cdot C = (\pi^*(H) - E_1 - E_2 - E_3 - E_4) \cdot C$$

gives $\pi^*(H) \cdot C = 2$, i.e. the image of C in the quintic model is a conic passing through two singular points a_i .

Lemma 5.2.

$$H_1(V, \mathbb{Z}) = 0.$$

Proof. It is enough to show that $\text{Tors } H_1(V, \mathbb{Z}) = 0$. Consider the bicanonical system $|2K_V|$ on V . A nonzero torsion divisor τ will cause a splitting

$$2K_V = (K_V + \tau) + (K_V - \tau).$$

For each nonzero torsion divisor τ the linear systems $|K_V + \tau|$ consists of a unique curve C_τ with $\chi(C_\tau, \mathcal{O}_{C_\tau}) = -1$. We have $C_\tau \cdot C_{-\tau} = 1$; moreover these two curves must intersect at a base point of $|3K_V|$ ([M], Theorem 2). Since $|3K_V - R|$ is base point free, this intersection point must lie on R , thus this member of $|2K|$ must intersect R at one point with multiplicity two.

Recall that the bicanonical system $|2K_V|$ is generated by the pullback of the pencil of quadrics $\mathcal{Q}(\lambda, \mu) = \lambda YT + \mu XZ$ in $\mathbb{P}^3$. We can also consider this system as the pullback of a pencil on V/σ^2 ; since $2K_V \cdot R = 2$, the general member of this pencil will intersect this rational curve at two points, and there will be exactly two members which intersect R at one point with multiplicity two. We have already two of them coming from the two reducible quadrics. In fact,

$$(\pi^*(H) - 2E_i - E_j - E_k) \cdot R = 1,$$

and σ^2 switches the two components of the reducible quadric. Thus any splitting of $2K_V$ as $C_\tau + C_{-\tau}$ must come from one of the reducible quadrics.

Let us consider these reducible members YT and XZ of the bicanonical system. Each hyperplane section X, Y, Z , and T of $\mathbb{P}^3$ is equivalent in V to $\pi^*(H) - 2E_i - E_j - E_k$ for the appropriate indices i, j, k . We have

$$(\pi^*(H) - 2E_i - E_j - E_k)^2 = -1$$

and

$$(\pi^*(H) - 2E_i - E_j - E_k) \cdot K_V = 1,$$

thus each irreducible component of the reducible quadrics defines on V a curve of arithmetic genus 1. Since $C_\tau^2 = C_\tau \cdot K_V = 1$, this shows that the curves C_τ cannot be irreducible.

Suppose C_τ is reducible. Since K_V is nef, and $C_\tau \cdot K_V = 1$, C_τ can contain only one irreducible component which is not a (-2) -curve. Let us denote this component by C'_τ . Let us write $C_\tau = C'_\tau + Z$, where Z is the sum of (-2) -curves. Since $C_\tau \cdot \pi^*(H) = K_V \cdot \pi^*(H) = 5$, we have $Z \cdot \pi^*(H) \leq 4$. From the previous lemma we know that for each irreducible component C of Z , we have $Z \cdot \pi^*(H) = 2$. Thus $Z = 0$, or $Z = C$ or $Z = C_1 + C_2$, where we do not assume that $C_1 \neq C_2$.

Case 1: $Z = C_1 + C_2$. We have

$$1 = K_V \cdot E_i = C_\tau \cdot E_i = C'_\tau \cdot E_i + (C_1 + C_2) \cdot E_i \geq (C_1 + C_2) \cdot E_i. \quad (5.1)$$

Recall that each (-2) -curve C intersects two E_i 's. If $C_1 = C_2$, we take E_i such that $C_1 \cdot E_i > 0$ and get a contradiction. If $C_1 \neq C_2$ and C_1, C_2 are not in the same divisor $D \in |3K_V - R|$, then we can find E_i which intersects both C_1 and C_2 . This follows from a simple combinatorial argument using the fact that each E_i intersects exactly one (-2) -component of a reducible divisor D from $|3K_V - R|$. So, applying (5.1), we get a contradiction.

If C_1 and C_2 are in the same divisor $D \in |3K_V - R|$, then

$$0 = K_V \cdot (C_1 + C_2) = C'_\tau \cdot (C_1 + C_2) - 4$$

implies that $C'_\tau \cdot (C_1 + C_2) = 4$; however this contradicts $C'_\tau \cdot (3K_V - R) = 2$.

Case 2: Z is irreducible. If Z is an irreducible (-2) -curve, then we have $K_V \cdot C'_\tau = 1$, $C'_\tau \cdot Z = 2$, $C'^2_\tau = -1$. Thus C'_τ is of arithmetic genus 1 and $C'_\tau \cdot Z = 2$. In the quintic model C'_τ is a plane cubic. The formula (5.1) shows that C'_τ intersects exactly

two E_i 's, say E_1 and E_2 , with multiplicity 1. The residual conic defines a divisor $Z' \in |\pi^*(H) - C'_\tau - aE_1 - bE_2|$. We have

$$Z'^2 = -2 + 2a + 2b - a^2 - b^2, \quad Z' \cdot K_V = 4 - a - b.$$

Since $Z'^2 + Z' \cdot K_V = -2$, we obtain

$$a(a-1) + b(b-1) = 4.$$

The only solution is $a = b = 2$. Thus

$$Z' \sim \pi^*(H) - C'_\tau - 2E_1 - 2E_2$$

and Z' is a (-2) -curve.

Since the image of C'_τ in the quintic model is a plane cubic, one of the plane components of the reducible quadric which contains $\pi(C'_\tau)$ must also contain the residual conic Z' , and

$$\pi^*(H) - 2E_i - E_j - E_k \sim C'_\tau + Z' \sim \pi^*(H) - 2E_1 - 2E_2.$$

This is obviously impossible.

Therefore a decomposition of $2K$ as $(K_V + \tau) + (K_V - \tau)$ cannot occur, and we have shown that there are no torsion divisors on V .

We next prove the following result:

Theorem 5.3.

$$\pi_1(V) = \{1\}.$$

By Lemma 5.2, it suffices to show that $\pi_1(V)$ is commutative.

As before, write $V' \rightarrow V$ for the blowup of the five isolated fixed points of σ^2 , and $p : V' \rightarrow F$ for the double cover of the rational surface F branched along the divisor $W' = \bar{W} + \sum_0^4 Z_i$ (see (4.3)). Recall that $R' = p^{-1}(\bar{W})$ is the pre-image of the curve R on V' , and $R_i = p^{-1}(Z_i)$ are the five (-1) -curves on V' blown up from the isolated fixed points of σ^2 . The curve $\bar{W}$ is fixed by σ^2 , while σ fixes Z_0 and two points on $\bar{W}$.

Since V' is birational to V , it suffices to show that $\pi_1(V')$ is commutative. The rational surface F is a blowup of the plane, so that $\pi_1(F - W')$ is generated by the image of the fundamental group of $L - W'$ where L is the pre-image on F of a line on $\mathbb{P}^2$. Abusing the terminology we shall still call L a line. Since V' is the double cover of F along W' , $\pi_1(V' - R' - \cup R_i)$ is a subgroup of index two in $\pi_1(F - W')$. The line L is isomorphic to $\mathbb{P}^1$, thus $\pi_1(L - W')$ is a free group on n generators, where $n = \#(L \cap W') - 1$. Generators for $\pi_1(L - W')$ can be chosen corresponding to small loops γ_i around the intersection points of L and W , such that the product $\gamma_0 \dots \gamma_n$ is trivial. The even products of the γ_i generate $\pi_1(V')$; the relations in $\pi_1(V')$ are $\gamma_i^2 = 1$ plus those coming from $\pi_1(F - W')$.

We will choose the line L on F by considering the fibres of the map $\varphi : V \rightarrow \mathbb{P}^1$ associated to the genus two pencil $|3K_V - R|$. This is the pullback of the pencil of lines in the plane through q , the order four point of the branch curve W . The member of this

pencil through Q_0 , the isolated fixed point of σ , is decomposed as $A_0 + A_0'$, the sum of two elliptic (-1) curves which are interchanged by σ . The curves A_0 and A_0' meet only at Q_0 , and each intersects R in three points. The pre-image of $A_0 + A_0'$ on V' is the σ -invariant divisor $A_0 + A_0' + R_0$; here we use the same notation for the proper transforms of A_0 , A_0' on V' .

Set $\ell = A_0/\sigma^2$; then $\sigma\ell = A_0'/\sigma^2$ and $\ell + \sigma\ell + Z_0$ is a σ -invariant fibre on F . Since σ switches the fibres of φ , this fibration also descends to $\bar{F} = F/\sigma$. On $\bar{F}$, ℓ and $\sigma\ell$ become identified. Set $\bar{\ell} = (\ell + \sigma\ell)/\sigma$ and $\bar{W}' = W'/\sigma$.

Lemma 5.4. ([KC], Theorem 2) *Let $Y \rightarrow \mathbb{P}^1$ be a fibration of a quasi-projective variety Y such that every fibre has a smooth component. Then for every fibre L the sequence*

$$\pi_1(L) \rightarrow \pi_1(Y) \rightarrow \pi_1(\mathbb{P}^1) = 1$$

is exact.

Thus we obtain a surjective homomorphism of groups

$$\pi_1(\ell + \sigma\ell - W') \rightarrow \pi_1(F - W').$$

Lemma 5.5. *The map*

$$\pi_1(\ell - W') \rightarrow \pi_1(F - W')$$

is surjective.

Proof. Consider the diagram

$$\begin{array}{ccc} \pi_1(\ell - W') & \rightarrow & \pi_1(F - W') \\ \downarrow & & \downarrow \\ \pi_1(\bar{\ell} - \bar{W}') & \rightarrow & \pi_1(\bar{F} - \bar{W}') \end{array}$$

By Lemma 5.4, the bottom horizontal map is surjective. Since $F - W'$ is an unramified cover of $\bar{F} - \bar{W}'$, $\pi_1(F - W') \rightarrow \pi_1(\bar{F} - \bar{W}')$ is injective. Also the left vertical map is surjective, since $\ell - W'$ is isomorphic to $\bar{\ell} - \bar{W}'$. Thus any element in $\pi_1(F - W')$ can be lifted to $\pi_1(\ell - W')$.

Thus the fundamental group of $F - W'$ is equal to the image of $\pi_1(\ell - W')$. Next consider the commutative diagram

$$\begin{array}{ccc} \pi_1(A_0 - R') & \rightarrow & \pi_1(V' - R' - \cup R_i) \\ \downarrow & & \downarrow \\ \pi_1(\ell - W') & \rightarrow & \pi_1(F - W'). \end{array}$$

The vertical arrows are injections onto a subgroup of index 2. By Lemma 5.5 the bottom horizontal arrow is surjective. Thus the top horizontal arrow is surjective. Finally consider the commutative diagram

$$\begin{array}{ccc} \pi_1(A_0 - R') & \rightarrow & \pi_1(V' - R' - \cup R_i) \\ \downarrow & & \downarrow \\ \pi_1(A_0) & \rightarrow & \pi_1(V') \end{array}$$

corresponding to the natural embeddings of the corresponding varieties. Since the vertical arrows are surjective, we obtain that the lower horizontal arrow is surjective. Thus the fundamental group of V' is the image of the fundamental group of an elliptic curve, which is commutative. Therefore $\pi_1(V) = H_1(V, \mathbb{Z}) = \{1\}$, and V is simply connected.

6. Comparison with Barlow surfaces

Here we prove that our surfaces are not isomorphic to Barlow surfaces. Since any Barlow surface contains four (-2) -curves it suffices to show that V does not contain any such curves.

Lemma 6.1. ([B], Proposition 1.3) *If S is a numerical Campedelli surface (i.e. $p_g = 0, K_S^2 = 2$), V is a simply connected surface, and $S \rightarrow V$ is a double cover branched along four (-2) -curves, then S has no even torsion.*

Theorem 6.2. *V contains no (-2) -curves and hence V is not isomorphic to a Barlow surface.*

Proof. By Lemma 5.1, the set of nodal curves consists of four disjoint curves $C, \sigma(C), \sigma^2(C)$, and $\sigma^3(C)$. We have

$$C + \sigma C + \sigma^2 C + \sigma^3 C \sim 2(3K_V - R) - 2A - 2A' = 2L,$$

where A, A' are two disjoint curves of arithmetic genus 1 with $A^2 = A'^2 = -1$.

We take the double cover of V branched along the four (-2) -curves, $f : Z \rightarrow V$. We have

$$L^2 = \frac{1}{4} \sum_0^3 (\sigma^i C)^2 = -2 \text{ and } L \cdot K_V = 0,$$

thus by Riemann-Roch, the Euler characteristic of Z is

$$\chi_Z = 2\chi_V + \frac{L \cdot (L + K_V)}{2} = 1.$$

We also have

$$K_Z^2 = 2(K_V + L)^2 = -2.$$

On Z the branch curves $\sigma^i C$ are disjoint (-1) -curves; blowing these four curves down gives a surface Z' with $K_{Z'}^2 = 2$. There are two possibilities for Z' : either $p_g = q = 1$, or $p_g = q = 0$, in which case Z' is a numerical Campedelli surface. We next show that neither case can occur, and therefore V cannot contain a (-2) -curve.

First suppose $p_g = q = 1$. Since $p_g(Z) = p_g(V) + h^0(V, \mathcal{O}_V(K_V + L))$ and $p_g(V) = 0$, the space $H^0(\mathcal{O}_V(K_V + L))$ is one-dimensional. We have

$$(K_V + L) \cdot \sigma^i(C) = L \cdot \sigma^i(C) = \frac{1}{2} \left(\sum_0^3 \sigma^i C \right) \cdot \sigma^i C = -1.$$

This implies that the unique divisor from $|K_V + L|$ contains all nodal curves $\sigma^i(C)$. But then $K_V + L - \sum_0^3 \sigma^i(C) = K_V + L - 2L = K_V - L$ is effective. We have $(K_V - L)^2 = -1$ and $K_V \cdot (K_V - L) = 1$. Thus $K_V - L$ is represented by a curve M of arithmetic genus 1. Also, from the proof of Lemma 5.1, each elliptic curve E_i intersects exactly two of the curves $\sigma^i(C)$. This implies $(K_V - L) \cdot E_i = 0$. If E_i is an irreducible component of the unique effective divisor $D \in |K_V - L|$, then all of the E_i 's are its irreducible components. Intersecting with K_V we see that it is impossible. Hence M is mapped isomorphically onto a curve $\bar{M}$ in the quintic model. Since, again by Lemma 5.1, $\pi^*(H) \cdot \sigma^i(C) = 2$ we have $\pi^*(H) \cdot (K_V - L) = 5 - 4 = 1$. Thus $\bar{M}$ is a line and cannot be of arithmetic genus 1. Therefore Z' cannot be a surface with $p_g = q = 1$.

Next consider the case $K_{Z'}^2 = 2$, $p_g = q = 0$. Then the minimal model Z' is a numerical Campedelli surface. By Corollary 3.6, the pencil $|3K - R|$ has five exceptional fibres which contribute to the Picard group 8 linearly independent divisors contained in fibres, namely the four (-2) -curves $C_1, \dots, C_4$ contained in two fibres, three disjoint (-1) -elliptic curves A_1, A_2, A_3 each contained in the other three exceptional fibres, plus the general fibre. It follows from the proof of Theorem 3.7 that each elliptic curve E_i intersects A_i or its complementary component A'_i with multiplicity 1. Also we know that each A_i is σ^2 -invariant. Thus we can choose the A_i in such a way that the elliptic curves E_1 and $E_2 = \sigma^2(E_1)$ intersect each A_i with multiplicity 1. The Picard number of V is nine. This implies that E_1 and E_2 (being 3-sections) differ by a $\mathbb{Q}$ -divisor contained in fibres. Write

$$E_1 = E_2 + n_1 A_1 + n_2 A_2 + n_3 A_3 + m_1 C_1 + m_2 C_2 + m_3 C_3 + m_4 C_4 + aF,$$

where F is a general fibre of $|3K - R|$. We also may assume that

$$C_2 = \sigma^2(C_1), C_4 = \sigma^2(C_3), E_1 \cdot C_1 = E_1 \cdot C_3 = 1, E_2 \cdot C_2 = E_2 \cdot C_4 = 1.$$

Intersecting both sides with the A_i and C_i gives

$$E_1 = E_2 - \frac{1}{2}C_1 - \frac{1}{2}C_2 + \frac{1}{2}C_3 + \frac{1}{2}C_4,$$

thus $2E_1 + C_1 + C_2 \sim 2E_2 + C_3 + C_4$ on V . On the double cover S of V , we have $2E'_1 + 2C'_1 + 2C'_2 \sim 2E'_2 + 2C'_3 + 2C'_4$ where we write E'_i, C'_i for the pullback of E_i, C_i to the cover. By Lemma 6.1, S has no 2-torsion, thus $E'_1 + C'_1 + C'_2 \sim E'_2 + C'_3 + C'_4$, and $|E'_1 + C'_1 + C'_2|$ is a genus two pencil on Z .

Now let A_1 be an elliptic component of another reducible fibre of $|3K_V - R|$; since A_1 does not meet the (-2) -curves C_i , it splits on the cover, and gives a section of the pencil $|E'_1 + C'_1 + C'_2|$ on Z . This implies that A_1 is an elliptic section of a rational pencil, a contradiction. Thus we cannot obtain such a covering of V , and again we see that there cannot be four (-2) -curves on V .

References

[B] Barlow, R., *A simply connected surface of general type with $p_g = 0$* , Invent. Math., 79 (1985), 293-301.

- [CG] Craighero, P., Gattazzo, R., *Quintic surfaces of $\mathbb{P}^3$ having a non singular model with $q = p_g = 0$, $P_2 \neq 0$* , Rend. Sem. Mat. Univ. Padova, 91 (1994), 187-198.
- [CL] Catanese, F., LeBrun, C., *On the scalar curvature of Einstein manifolds*, preprint.
- [KC] Keum, J.H., Chung, I.J., *On fundamental groups of fibred complex manifolds*, Mich. Math. Journ., to appear.
- [M] Miyaoka, Y., *Tricanonical maps of numerical Godeaux surfaces*, Invent. math., 34 (1976) , 99-111.
- [OP] Oort, F., Peters, C., *A Campedelli surface with torsion group $\mathbb{Z}/2\mathbb{Z}$* , Indag. Math., 43 (1981), 399-407.
- [R] Reid, M., *Campedelli versus Godeaux*, in “Problems in the theory of surfaces and their classification (Cortona, 1988)”, pp. 309-365, Academic Press, 1991.
- [St] Stagnaro, E., *On Campedelli branch loci*, Dip. di Met. e Mod. Matematici, Università di Padova, 53, (1996).
- [W] Werner, C., *A surface of general type with $p_g = q = 0$, $K^2 = 1$* , Manuscripta math., 84 (1994), 327-341.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109
E-mail address: idolga@math.lsa.umich.edu, cwerner@math.lsa.umich.edu